I want to be able	to:			
My asthma Patient name:	Children's Healthcare of Atlanta Dedicated to All Botton			
Doctor's name Doctor's phone #:		Signature		
Controller medicines	How much to take	How often	Other instructions	
		times per day EVERY DAY  times per day EVERY DAY  times per day EVERY DAY	্ৰ Gargle or rinse mouth after use	
Quick-relief medicines	How much to take	How often	Other instructions	
	☐ 2 puffs ☐ 4-6 puffs ☐ 1 nebulizer treatment	Take ONLY as needed (see below — starting in Yellow Zone or before excercise)	NOTE: If you need this medicine more than 2 days a week, call your doctor.	
Asthma triggers (check all t Exercise Change in Pollens Respirator	temperature 🖵 Molds	☐ Strong emotions	odors or fumes	
Doing well.     No coughing, wheezing tightness, shortness of buring the day or night     Can go to school and place.	preath	PREVENT asth  Take my cor  Before exer	ma symptoms every day:  ntroller medicines (above) every day  cise, take puff(s) of  ers that make my asthma worse	
Be careful.  Coughing, wheezing, che tightness, shortness of be Waking at hight due to asthma symptoms  Can do some, but not all usual activities  Runny nose, watery eyes	oreath	CAUTION. Continue taking my controller medicines every day  Take puffs or nebulizer treatment(s) of quick relief medicine. If I am not back in the Green Zone within one hour, then I should:  Continue using quick relief medicine every 4 hours as needed. Call provider if not improving in days.  Increase Add		
Ask for help.	(۲۶٪	MEDICAL ALE	RT! Get help!	



- Very short of breath
- Continual coughing
- Škin between ribs is pulling inwards
- Difficulty speaking without running out of breath
- Quick-relief medicines have not helped
- Symptoms same or worse after 48 hours in Yellow Zone



Take quick-relief medicine: \_\_\_\_ puffs every \_\_\_ minutes and get help immediately.

 Take	A STATE OF THE STA

	Ċall			

If skin, fingernail or lip color is blue at any time:

Call 911 for help or go to the nearest Emergency Department