Margaret Mace School 1201 Atlantic Avenue, North Wildwood, NJ 08260 (609) 522-1454 fax (609)846-1704

Dear Parent/Guardian,

To participate in school sports, there are a few health related forms that need to be completed and returned **PRIOR** to the student's first (1^{st}) practice/tryout. The list below contains the requirements:

To be completed **PRIOR** to first (1^{st}) practice/tryout:

- Physical Evaluation (YELLOW forms completed by physician)
- ALL WHITE forms need to be reviewed by Parent/Guardian and student

The following forms require **signatures** from **BOTH** Parent/Guardian and student.

- 1. Health History
- 2. Sports-Related Concussion and Head Injury (signature required on BACK)
- 3. Sudden Cardiac Death sign-off sheet
- 4. Use and Misuse of Opioids

Health History Update Form- If your child has a current completed physical packet (within 365 days) on file in the nurse's office, only the Health History Update Form is needed at the beginning of each sports season.

ALL forms are listed on the Nurse and Athletics sections of the school's website. Additionally, there are packets available in the nurse's office.

Per state law, ALL requirements need to be completed **PRIOR** to first practice/tryout of each sport.

Thank you, Renee M. Forrest RN, BSN,CSN School nurse (ext 617)

NOTE: The preparticiaption physical examination must be conducted by a health care provider who 1) is a licensed physician, advanced practice nurse, or physician assistant; and 2) completed the Student-Athlete Cardiac Assessment Professional Development Module.

Date of birth

PREPARTICIPATION PHYSICAL EVALUATION PHYSICAL EXAMINATION FORM

Name

PHYSICIAN REMINDERS

1. Consider additional questions on more sensitive issues

- Do you feel stressed out or under a lot of pressure? Do you ever feel sad, hopeless, depressed, or anxious? .
- * Do you feel safe at your home or residence?
- * Have you ever tried cigarettes, chewing tobacco, snuff, or dip?
- During the past 30 days, did you use chewing tobacco, snuff, or dip?
- Do you drink alcohol or use any other drugs?
- Have you ever taken anabolic steroids or used any other performance supplement?
- * Have you ever taken any supplements to help you gain or lose weight or improve your performance?
- Do you wear a seat belt, use a helmet, and use condoms? 2. Consider reviewing questions on cardiovascular symptoms (questions 5-14).

EXAMINATION

	INATION	1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1		and all and							1000 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100	A LANGER	and the second	
Height			١	Neight		🗆 Ma	ale 🗆	I Female						
BP	1	(1)	Pulse	Visi	on R 20)/	L 20/		Corrected E	1 Y		
MEDIC	CAL							NORMAL	1123	AB	NORMAL FINDI	NGS	A MARTINESS	
Appear Mar arm	rance rfan stigmata (k 1 span > height,	yphoscoliosis hyperlaxity, r	, high-arc nyopia, M	hed palate	e, pectus e insufficien	excavatum, arachnodactyly, icy)								
	ars/nose/throat ills equal iring													
Lymph	nodes													
	rmurs (ausculta ation of point of				a)									
PulsesSim	ultaneous femo	ral and radial	pulses											
Lungs														
Abdom	en													
Genitou	urinary (males c	nly)⁵												
	, lesions sugges	stive of MRSA	, tinea coi	rporis										
Neurolo														
	ULOSKELETAL	E. 74.2						「「ないた」の目的	Les and					
Neck														
Back			-											
Should							-							
Elbow/	and the second se													
	and/fingers											_		
Hip/thig	gh 🛛 👘													
Knee														
Leg/anl														
Foot/toe	and the second se													_
Functio	nal													

Duck-walk, single leg hop

*Consider ECG, echocardiogram, and referral to cardiology for abnormal cardiac history or exam.

*Consider GU exam if in private setting. Having third party present is recommended. *Consider cognitive evaluation or baseline neuropsychiatric testing if a history of significant concussion.

Cleared for all sports without restriction

-	cleared for all sports without restriction with re	
	Not cleared	
	Pending further evaluation	
	For any sports	
	For certain sports	
	Reason	
Rei	commendations	

I have examined the above-named student and completed the preparticipation physical evaluation. The athlete does not present apparent clinical contraindications to practice and participate in the sport(s) as outlined above. A copy of the physical exam is on record in my office and can be made available to the school at the request of the parents. If conditions arise after the athlete has been cleared for participation, a physician may rescind the clearance until the problem is resolved and the potential consequences are completely explained to the athlete (and parents/guardians).

Name of physician, advanced practice nurse (APN), physician assistant (PA) (print/type)	Date of exam
Address	Phone
Signature of physician, APN, PA	
-3	

©2010 American Academy of Family Physicians, American Academy of Pediatrics, American College of Sports Medicine, American Medical Society for Sports Medicine, American Orthopaedic Society for Sports Medicine, and American Osteopathic Academy of Sports Medicine. Permission is granted to reprint for noncommercial, educational purposes with acknowledgment. HE0503 9-2681/0410

New Jersey Department of Education 2014; Pursuant to P.L.2013, c.71

PREPARTICIPATION PHYSICAL EVALUATION CLEARANCE FORM

Name	_ Sex 🗆 M 🗆 F Age	Date of birth	The second
Cleared for all sports without restriction			
Cleared for all sports without restriction with recommendations for further even	valuation or treatment for		
□ Not cleared			
Pending further evaluation			
 For any sports 			
□ For certain sports			
Reason			
Recommendations			
Allergies			(States
Other information			
Other information			
HCP OFFICE STAMP	SCHOOL PHYSICIAN:		
	Reviewed on		
		(Date)	
	Approved Not	Approved	
	Signature:		
I have examined the above-named student and completed the prep clinical contraindications to practice and participate in the sport(s) and can be made available to the school at the request of the paren the physician may rescind the clearance until the problem is resolv (and parents/guardians).	as outlined above. A copy of the its. If conditions arise after the at	physical exam is on record i hlete has been cleared for pa	n my office articipation,
Name of physician, advanced practice nurse (APN), physician assistant (PA)		Date	
Address		Phone	

Signature of physician, APN, PA_

Completed Cardiac Assessment Professional Development Module

_ Signature_

Date_____

© 2010 American Academy of Family Physicians, American Academy of Pediatrics, American College of Sports Medicine, American Medical Society for Sports Medicine, American Orthopaedic Society for Sports Medicine, and American Osteopathic Academy of Sports Medicine. Permission is granted to reprint for noncommercial, educational purposes with acknowledgment. New Jersey Department of Education 2014; Pursuant to P.L.2013, c.71

PREPARTICIPATION PHYSICAL EVALUATION THE ATHLETE WITH SPECIAL NEEDS: SUPPLEMENTAL HISTORY FORM

Date of Exa	n					
Name				Date of birth		
Sex	Age	Grade	School	Sport(s)		ana ana ang ang ang ang ang ang ang ang
1. Type of	disability					
2. Date of	disability					
3. Classific	ation (if available)					
4. Cause of	disability (birth, dis	ease, accident/trauma, other)			
5. List the	sports you are intere	sted in playing				
					Yes	No
6. Do you r	egularly use a brace	, assistive device, or prosthe	tic?			
7. Do you u	se any special brac	e or assistive device for spor	ts?			
8. Do you h	ave any rashes, pre	ssure sores, or any other ski	n problems?			
9. Do you h	ave a hearing loss?	Do you use a hearing aid?				
10. Do you h	ave a visual impairr	nent?				
11. Do you u	se any special devic	es for bowel or bladder func	tion?			
12. Do you h	ave burning or disco	omfort when urinating?				
13. Have you	I had autonomic dys	reflexia?				
14. Have you	ever been diagnos	ed with a heat-related (hyper	thermia) or cold-related (hypothermia) illn	ess?		
15. Do you h	ave muscle spastici	ty?				
16. Do you h	ave frequent seizure	es that cannot be controlled l	by medication?			

Explain "yes" answers here

	Y	es No
Atlantoaxial instability		
X-ray evaluation for atlantoaxial instability		
Dislocated joints (more than one)		
Easy bleeding		
Enlarged spleen		
Hepatitis		
Osteopenia or osteoporosis		
Difficulty controlling bowel		
Difficulty controlling bladder		
Numbness or tingling in arms or hands		
Numbness or tingling in legs or feet		
Weakness in arms or hands		
Weakness in legs or feet		
Recent change in coordination	1	
Recent change in ability to walk		
Spina bifida		
Latex allergy		

Explain "yes" answers here

I hereby state that, to the best of my knowledge, my answers to the above questions are complete and correct.

Signature of athlete

____ Signature of parent/guardian _____

©2010 American Academy of Family Physicians, American Academy of Pediatrics, American College of Sports Medicine, American Medical Society for Sports Medicine, American Orthopaedic Society for Sports Medicine, and American Osteopathic Academy of Sports Medicine. Permission is granted to reprint for noncommercial, educational purposes with acknowledgment. New Jersey Department of Education 2014; Pursuant to P.L.2013, c.71

Date_

PREPARTICIPATION PHYSICAL EVALUATION HISTORY FORM

Name				Date of birth	τΰ.	6
				Sport(s)		
Medicines and All	ergies: Please list all of the prescription and ov	er-the-c	ounter	medicines and supplements (herbal and nutritional) that you are currently	taking	
			Journes	menones and subbenence (ternar and induced) mar for a contently	aning	
Do you have any all	lergies? 🗆 Yes 🖾 No lf yes, please ic	lentify s	pecific	allergy below.		
🗆 Medicines	□ Pollens			□ Food □ Stinging Insects		
xolain "Yes" answe	rs below. Circle questions you don't know the a	mswers	tn.			
GENERAL QUESTIONS	the second s	1 Ses	-	Medical questions	Ves	
1. Has a doctor ever any reason?	denied or restricted your participation in sports for			26. Do you cough, wheeze, or have difficulty breathing during or after exercise?		
2. Do you have any o	ngoing medical conditions? If so, please identify		1	27. Have you ever used an inhaler or taken asthma medicine?		
	a 🖾 Anemia 🖾 Diabetes 🖾 Infections			28. Is there anyone in your family who has asthma?		
Other: 3. Have you ever spen	nt the night in the hospital?			29. Were you born without or are you missing a kidney, an eye, a testicle (males), your spleen, or any other organ?		
4. Have you ever had				30. Do you have groin pain or a painful bulge or hernía in the groin area?		
IEART REALTY OBES	A COMPANY OF THE DESIGN OF THE OTHER DESIGN OF THE	Ves	No	31. Have you had infectious mononucleosis (mono) within the last month?		
Have you ever pass AFTER exercise?	sed out or nearly passed out DURING or			32. Do you have any rashes, pressure sores, or other skin problems?		
	discomfort, pain, tightness, or pressure in your			33. Have you had a herpes or MRSA skin infection?		
chest during exerci				34. Have you ever had a head injury or concussion?		L
7. Does your heart ev	er race or sldp beats (irregular beats) during exercise?			35. Have you ever had a hit or blow to the head that caused confusion, prolonged headache, or memory problems?		
8. Has a doctor ever to	old you that you have any heart problems? If so,			36. Do you have a history of seizure disorder?		
check all that apply High blood pres			1	37. Do you have headaches with exercise?		
High cholestero	A heart infection			38. Have you ever had numbness, tingling, or weakness in your arms or legs after being hit or falling?		
9. Has a doctor ever o echocardiogram)	rdered a test for your heart? (For example, ECG/EKG,			39. Have you ever been unable to move your arms or legs after being hit or falling?		
0. Do you get lighthead	ded or feel more short of breath than expected	1	1	40. Have you ever become ill while exercising in the heat?		
during exercise?			ļ	41. Do you get frequent muscle cramps when exercising?		
	an unexplained seizure?			42. Do you or someone in your family have sickle cell trait or disease?		
 Do you get more tire during exercise? 	ed or short of breath more quickly than your friends			43. Have you had any problems with your eyes or vision?		_
Traper and the second s	ions about your family	Ves	Ne	44. Have you had any eye injuries?		_
A A A A A A A A A A A A A A A A A A A	ber or relative died of heart problems or had an	100		45. Do you wear glasses or contact lenses?		
unexpected or unexp	plained sudden death before age 50 (including			46. Do you wear protective eyewear, such as goggles or a face shield?		
. Does anyone in your	ed car accident, or sudden infant death syndrome)? family have hypertrophic cardiomyopathy, Marfan			47. Do you worry about your weight?48. Are you trying to or has anyone recommended that you gain or loss weight?		
syndrome, short QT :	ogenic right ventricular cardiomyopathy, long QT syndrome, Brugada syndrome, or catecholaminergic			lose weight? 49. Are you on a special diet or do you avoid certain types of foods?		
polymorphic ventricu				50. Have you ever had an eating disorder?		
 Does anyone in your implanted defibrillato 	family have a heart problem, pacemaker, or			51. Do you have any concerns that you would like to discuss with a doctor?		
	amily had unexplained fainting, unexplained			FEMALES ONLY		53
seizures, or near dron				52. Have you ever had a menstrual period?		-
NE AND JOINT QUES	TIONS	Ves	lio	53. How old were you when you had your first menstrual period?		
	i injury to a bone, muscle, ligament, or tendon iss a practice or a game?			54. How many periods have you had in the last 12 months?		
Have you ever had an	y broken or fractured bones or dislocated joints?			explain "yes" answers here		
	injury that required x-rays, MRI, CT scan, brace, a cast, or crutches?					

21. Have you ever been told that you have or have you had an x-ray for neck instability or atlantoaxial instability? (Down syndrome or dwarfism) 22. Do you regularly use a brace, orthotics, or other assistive device? 23. Do you have a bone, muscle, or joint injury that bothers you? 24. Do any of your joints become painful, swollen, feel warm, or look red? 25. Do you have any history of juvenile arthritis or connective tissue disease? I hereby state that, to the best of my knowledge, my answers to the above questions are complete and correct.

Signature of athlete

20, Have you ever had a stress fracture?

4

Q

Date

©2010 American Academy of Family Physicians, American Academy of Pediatrics, American College of Sports Medicine, American Medical Society for Sports Medicine, American Orthopæedic Society for Sports Medicine, and American Osteopathic Academy of Sports Medicine. Permission is granted to reprint for noncommercial, educational purposes with acknowledgment. HEOSOS

Signature of parent/guardian

New Jersey Department of Education 2014; Pursuant to P.L.2013, c.71

State of New Jersey Department of Education <u>HEALTH HISTORY UPDATE QUESTIONNAIRE</u>

Name	of	School	
TIMIT	OT.	OCHOOL	

	•	Age	Gr	ade
tudentate of Last Physical Examination	Sport	1,±5°_		
nce the last pre-participation physical examination, has your son/o	daughter:			
 Been medically advised not to participate in a sport? If yes, describe in detail 		Yes	No	
		·····		
2 Sustained a concussion been unconscious or last and the second		· ·		
2. Sustained a concussion, been unconscious or lost memory from a b	low to the he	di Ves	No	
If yes, explain in detail		iu! ies	 · · · ·	
			•	
	· · · · ·		,	
3. Broken a bone or sprained/strained/dislocated any muscle or joints?	2	Yes	No	•
If yes, describe in detail		· · ·		
4. Fainted or "blacked out?" If yes, was this during or immediately after exercise?		. Van	Mo	
	•			
. Experienced chest pains, shortness of breath or "racing heart?"	8	Yes	_ No	
. Has there been a recent history of fatigue and unusual tiredness?		Yes	No	
. Been hospitalized or had to go to the emergency room?		Yes		
If yes, explain in detail		. 105	NO	
		• :		• •
			·.	
Since the last physical examination, has there been a sudden death in under age 50 had a heart attack or "heart trouble?"	the family or	has any me Yes	mber of the	e family
Started or stopped taking any over-the-counter or prescribed medicati	ions?		No .	۰.
If yes, name of medication(s)			· · · ·	· . ·

Please sign Back Page

PARENT & ATHLETE CONCUSSION INFORMATION SHEET

WHAT IS A CONCUSSION?

A concussion is a type of traumatic brain injury that changes the way the brain normally works. A concussion is caused by a bump, blow, or jolt to the head or body that causes the head and brain to move quickly back and forth. Even a "ding," "getting your bell rung," or what seems to be a mild bump or blow to the head can be serious.

WHAT ARE THE SIGNS AND SYMPTOMS OF CONCUSSION?

Signs and symptoms of concussion can show up right after the injury or may not appear or be noticed until days or weeks after the injury.

If an athlete reports one or more symptoms of concussion after a bump, blow, or jolt to the head or body, s/he should be kept out of play the day of the injury. The athlete should only return to play with permission from a health care professional experienced in evaluating for concussion.

DID YOU KNOW?

- Most concussions occur without loss of consciousness.
- Athletes who have, at any point in their lives, had a concussion have an increased risk for another concussion.
- Young children and teens are more likely to get a concussion and take longer to recover than adults.

SYMPTOMS REPORTED BY ATHLETE:

- · Headache or "pressure" in head
- Nausea or vomiting
- · Balance problems or dizziness
- Double or blurry vision
- Sensitivity to light
- Sensitivity to noise
- Feeling sluggish, hazy, foggy, or groggy
- · Concentration or memory problems
- Confusion
- Just not "feeling right" or is "feeling down"

SIGNS OBSERVED BY COACHING STAFF:

- Appears dazed or stunned
- Is confused about assignment or position
- · Forgets an instruction
- Is unsure of game, score, or opponent
- Moves clumsily
- Answers questions slowly
- Loses consciousness (even briefly)
- · Shows mood, behavior, or personality changes
- · Can't recall events prior to hit or fall
- · Can't recall events after hit or fall

"IT'S BETTER TO MISS ONE GAME THAN THE WHOLE SEASON"

CONCUSSION DANGER SIGNS

In rare cases, a dangerous blood clot may form on the brain in a person with a concussion and crowd the brain against the skull. An athlete should receive immediate medical attention if after a bump, blow, or jolt to the head or body s/he exhibits any of the following danger signs:

- · One pupil larger than the other
- · Is drowsy or cannot be awakened
- A headache that gets worse
- · Weakness, numbness, or decreased coordination
- · Repeated vomiting or nausea
- Slurred speech
- · Convulsions or seizures
- · Cannot recognize people or places
- · Becomes increasingly confused, restless, or agitated
- · Has unusual behavior
- Loses consciousness (even a brief loss of consciousness should be taken seriously)

WHAT SHOULD YOU DO IF YOU THINK YOUR ATHLETE HAS A CONCUSSION?

- If you suspect that an athlete has a concussion, remove the athlete from play and seek medical attention. Do not try to judge the severity of the injury yourself. Keep the athlete out of play the day of the injury and until a health care professional, experienced in evaluating for concussion, says s/he is symptom-free and it's OK to return to play.
- 2. Rest is key to helping an athlete recover from a concussion. Exercising or activities that involve a lot of concentration, such as studying, working on the computer, and playing video games, may cause concussion symptoms to reappear or get worse. After a concussion, returning to sports and school is a gradual process that should be carefully managed and monitored by a health care professional.
- 3. Remember: Concussions affect people differently. While most athletes with a concussion recover quickly and fully, some will have symptoms that last for days, or even weeks. A more serious concussion can last for months or longer.

WHY SHOULD AN ATHLETE REPORT THEIR SYMPTOMS?

If an athlete has a concussion, his/her brain needs time to heal. While an athlete's brain is still healing, s/he is much more likely to have another concussion. Repeat concussions can increase the time it takes to recover. In rare cases, repeat concussions in young athletes can result in brain swelling or permanent damage to their brain. They can even be fatal.

STUDENT-ATHLETE NAME PRINTED

STUDENT-ATHLETE NAME SIGNED

DATE

PARENT OR GUARDIAN NAME PRINTED

PARENT OR GUARDIAN NAME SIGNED

DATE

JOIN THE CONVERSATION L www.facebook.com/CDCHeadsUp

HEADS UP

TO LEARN MORE GO TO >> WWW.CDC.GOV/CONCUSSION

Content Source: CDC's Heads Up Program. Created through a grant to the CDC Foundation from the National Operating Committee on Standards for Athletic Equipment (NOCSAE).

OPIOID USE AND MISUSE EDUCATIONAL FACT SHEET Keeping Student-Athletes Safe

School athletics can serve an integral role in students' development. In addition to providing healthy forms of exercise, school athletics foster friendships and camaraderie, promote sportsmanship and fair play, and instill the value of competition.

Unfortunately, sports activities may also lead to injury and, in rare cases, result in pain that is severe or long-lasting enough to require a prescription opioid painkiller.¹ It is important to understand that overdoses from opioids are on the rise and are killing Americans of all ages and backgrounds. Families and communities across the country are coping with the health, emotional and economic effects of this epidemic.²

This educational fact sheet, created by the New Jersey Department of Education as required by state law (*N.J.S.A.* 18A:40-41.10), provides information concerning the use and misuse of opioid drugs in the event that a health care provider prescribes a student-athlete or cheerleader an opioid for a sports-related injury. Student-athletes and cheerleaders participating in an interscholastic sports program (and their parent or guardian, if the student is under age 18) must provide their school district written acknowledgment of their receipt of this fact sheet.

How Do Athletes Obtain Opioids?

In some cases, student-athletes are prescribed these medications. According to research, about a third of young people studied obtained pills from their own previous prescriptions (i.e., an unfinished prescription used outside of a physician's supervision), and 83 percent of adolescents had unsupervised access to their prescription medications.³ It is important for parents to understand the possible hazard of having unsecured prescription medications in their households. Parents should also understand the importance of proper storage and disposal of medications, even if they believe their child would not engage in non-medical use or diversion of prescription medications.

What Are Signs of Opioid Use?

According to the National Council on Alcoholism and Drug Dependence, 12 percent of male athletes and 8 percent of female athletes had used prescription opioids in the 12-month period studied.³ In the early stages of abuse, the athlete may exhibit unprovoked nausea and/or vomiting. However, as he or she develops a tolerance to the drug, those signs will diminish. Constipation is not uncommon, but may not be reported. One of the most significant indications of a possible opioid addiction is an athlete's decrease in academic or athletic performance, or a lack of interest in his or her sport. If these warning signs are noticed, best practices call for the student to be referred to the appropriate professional for screening,⁴ such as provided through an evidence-based practice to identify problematic use, abuse and dependence on illicit drugs (e.g., Screening, Brief Intervention, and Referral to Treatment (SBIRT)) offered through the New Jersey Department of Health.

What Are Some Ways Opioid Use and Misuse Can Be Prevented?

According to NJSIAA Sports Medical Advisory Committee chair, John P. Kripsak, D.O., "Studies indicate that about 80 percent of heroin users started out by abusing narcotic painkillers."

According to the New Jersey State Interscholastic Athletic Association (NJSIAA) Sports Medical Advisory Committee chair, John P. Kripsak, D.O., "Studies indicate that about 80 percent of heroin users started out by abusing narcotic painkillers."

The Sports Medical Advisory Committee, which includes representatives of NJSIAA member schools as well as experts in the field of healthcare and medicine, recommends the following:

- The pain from most sports-related injuries can be managed with non-narcotic medications such as acetaminophen, nonsteroidal anti-inflammatory medications like ibuprofen, naproxen or aspirin. Read the label carefully and always take the recommended dose, or follow your doctor's instructions. More is not necessarily better when taking an over-the-counter (OTC) pain medication, and it can lead to dangerous side effects.⁴
- Ice therapy can be utilized appropriately as an anesthetic.
- Always discuss with your physician exactly what is being prescribed for pain and request to avoid narcotics.
- In extreme cases, such as severe trauma or post-surgical pain, opioid pain medication should not be prescribed for more than five days at a time;
- Parents or guardians should always control the dispensing of pain medications and keep them in a safe, non-accessible location; and
- Unused medications should be disposed of immediately upon cessation of use. Ask your pharmacist about drop-off locations
 or home disposal kits like Deterra or Medsaway.



STATE OF NEW JERSEY DEPARTMENT OF EDUCATION

In consultation with NJSIAA SPORTS MEDICAL

ADVISORY COMMITTEE

Karan Chauhan Parsippany Hills High School, **Permanent Student Representative** New Jersey State Board of Education



STATE OF NEW JERSEY

Even With Proper Training and Prevention, **Sports Injuries May Occur**

There are two kinds of sports injuries. Acute injuries happen suddenly, such as a sprained ankle or strained back. Chronic injuries may happen after someone plays a sport or exercises over a long period of time, even when applying overuse-preventative techniques.⁵

Athletes should be encouraged to speak up about injuries, coaches should be supported in injury-prevention decisions, and parents and young athletes are encouraged to become better educated about sports safety.⁶

What Are Some Ways to Reduce the Risk of Injury?'

Half of all sports medicine injuries in children and teens are from overuse. An overuse injury is damage to a bone, muscle, ligament, or tendon caused by repetitive stress without allowing time for the body to heal. Children and teens are at increased risk for overuse injuries because growing bones are less resilient to stress. Also, young athletes may not know that certain symptoms are signs of overuse.

The best way to deal with sports injuries is to keep them from happening in the first place. Here are some recommendations to consider:



NJ Health

PREPARE Obtain the preparticipation physical evaluation prior to participation on a school-sponsored interscholastic or intramural athletic team or squad.



TRAINING Increase weekly training time, mileage or repetitions no more than 10 percent per week. For example, if running 10 miles one week, increase to 11 miles the following week. Athletes should also cross-train and perform sport-specific drills in different ways, such as running in a swimming pool instead of only running on the road.



CONDITIONING Maintain a good fitness level during the season and offseason. Also important are proper warm-up and cooldown exercises.



ADEQUATE HYDRATION Keep the body hydrated to help the heart more easily pump blood to muscles, which helps muscles work efficiently.



REST UP Take at least one day off per week from organized activity to recover physically and mentally. Athletes should take a combined three months off per year from a specific sport (may be divided throughout the year in one-month increments). Athletes may remain physically active during rest periods through alternative low-stress activities such as stretching, yoga or walking.



PROPER EQUIPMENT Wear appropriate and properly fitted protective equipment such as pads (neck, shoulder, elbow, chest, knee, and shin), helmets, mouthpieces, face guards, protective cups, and eyewear. Do not assume that protective gear will prevent all injuries while performing more dangerous or risky activities.

Resources for Parents and Students on Preventing Substance Misuse and Abuse

The following list provides some examples of resources:

National Council on Alcoholism and Drug Dependence - NJ promotes addiction treatment and recovery.

New Jersey Department of Health, Division of Mental Health and Addiction Services is committed to providing consumers and families with a wellness and recovery-oriented model of care.

New Jersey Prevention Network includes a parent's quiz on the effects of opioids.

Operation Prevention Parent Toolkit is designed to help parents learn more about the opioid epidemic, recognize warning signs, and open lines of communication with their children and those in the community.

Parent to Parent NJ is a grassroots coalition for families and children struggling with alcohol and drug addiction.

Partnership for a Drug Free New Jersey is New Jersey's anti-drug alliance created to localize and strengthen drug-prevention media efforts to prevent unlawful drug use, especially among young people.

The Science of Addiction: The Stories of Teens shares common misconceptions about opioids through the voices of teens.

Youth IMPACTing NJ is made up of youth representatives from coalitions across the state of New Jersey who have been impacting their communities and peers by spreading the word about the dangers of underage drinking, marijuana use, and other substance misuse.

References¹ Massachusetts Technical Assistance Partnership for Prevention

- Committee (SMAC)
- ² Centers for Disease Control and Prevention
- Association (NJSIAA) Sports Medical Advisory ⁴ Athletic Management, David Csillan, athletic
 - and Skin Diseases ⁶ USA TODAY

³ New Jersey State Interscholastic Athletic

trainer, Ewing High School, NJSIAA SMAC

⁷ American Academy of Pediatrics

⁵ National Institute of Arthritis and Musculoskeletal

An online version of this fact sheet is available on the New Jersey Department of Education's Alcohol, Tobacco, and Other Drug Use webpage. Updated Jan. 30, 2018.

North Wildwood School District Margaret Mace School Office of the Superintendent 1201 Atlantic Avenue North Wildwood, New Jersey 08260 carmstrong@mmace.com

Administration: (609) 522-6885

Fax: (609) 522-2308 Brenda G. Trasatti Secretary to the Superintendent

School: (609) 522-1454 Christopher Armstrong Superintendent

ið 1

Use and Misuse of Opioid Drugs Fact Sheet Student-Athlete and Parent/Guardian Sign-Off

In accordance with *N.J.S.A. 18A:40-41.10*, public school districts, approved private schools for students with disabilities, and nonpublic schools participating in an interscholastic sports program must distribute this *Opioid Use and Misuse Educational Fact Sheet* to all students-athletes and cheerleaders. In addition, schools and districts must obtain a signed acknowledgement of receipt of the fact sheet from each student-athlete and cheerleader, and for students under 18, the parent or guardian must also sign.

This sign-off sheet is due to the Head Coach of your sport prior to the first official practice of the 2018/2019 school year as determined by the New Jersey State Interscholastic Athletic Association and is required annually prior to the student-athlete's or cheerleader's first official practice of the school year.

Name of School: Margaret Mace School

Name of School District: North Wildwood

I/We acknowledge that we received and reviewed the Educational Fact Sheet on the Use and Misuse of Opioid Drugs.

Student Signature:

Date:

*Does not include athletic clubs or intramural events.

SPORTS-RELATED EYE INJURIES:

AN EDUCATIONAL FACT SHEET FOR PARENTS

Participating in sports and recreational activities is an important part of a healthy, physically active lifestyle for children. Unfortunately, injuries can, and do, occur. Children are at particular risk for sustaining a sports-related eye injury and most of these injuries can be prevented. Every year, more than 30,000 children sustain serious sports-related eye injuries. Every 13 minutes, an emergency room in the United States treats a sports-related eye injury.¹ According to the National Eye Institute, the sports with the highest rate of eye injuries are: baseball/softball, ice hockey, racquet sports, and basketball, followed by fencing, lacrosse, paintball and boxing.

Thankfully, there are steps that parents can take to ensure their children's safety on the field, the court, or wherever they play or participate in sports and recreational activities.

Prevention of Sports-Related Eye Injuries

Approximately 90% of sports-related eye injuries can be prevented with simple precautions, such as using protective eyewear.² Each sport has a certain type of recommended protective eyewear, as determined by the American Society for Testing and Materials (ASTM). Protective eyewear should sit comfortably on the face. Poorly fitted equipment may be uncomfortable, and may not offer the best eye protection. Protective eyewear for sports includes, among other things, safety goggles and eye guards, and it should be made of polycarbonate lenses, a strong, shatterproof plastic. Polycarbonate lenses are much stronger than regular lenses.³

Health care providers (HCP), including family physicians, ophthalmologists, optometrists, and others, play a critical role in advising students, parents and guardians about the proper use

of protective eyewear. To find out what kind of eye protection is recommended, and permitted for your child's sport, visit the National Eye Institute at http://www.nei.nih.gov/sports/findingprotection.asp. Prevent Blindness America also offers tips for choosing and buying protective eyewear at http://www.preventblindness.org/tips-buying-sports-eye-protectors, and http://www.preventblindness.org/ recommended-sports-eye-protectors.

It is recommended that all children participating in school sports or recreational sports wear protective eyewear. Parents and coaches need to make sure young athletes protect their eyes, and properly gear up for the game. Protective eyewear should be part of any uniform to help reduce the occurrence of sports-related eye injuries. Since many youth teams do not require eye protection, parents may need to ensure that their children wear safety glasses or goggles whenever they play sports. Parents can set a good example by wearing protective eyewear when they play sports.

- ¹ National Eye Institute, National Eye Health Education Program, Sports-Related Eye Injuries: What You Need to Know and Tips for Prevention, www.nei.nih.gov/sports/pdf/sportsrelatedeyeInjuries.pdf, December 26, 2013.
- ² Rodriguez, Jorge O., D.O., and Lavina, Adrian M., M.D., Prevention and Treatment of Common Eye Injuries in Sports, http://www.aafp.org/afp/2003/0401/p1481.html, September 4, 2014; National Eye Health Education Program, Sports-Related Eye Injuries: What You Need to Know and Tips for Prevention, www.nei.nih.gov/sports/pdf/sportsrelatedeyeInjuries.pdf, December 26, 2013.
- ³ Bedinghaus, Troy, O.D., Sports Eye Injuries, http://vision.about.com/od/emergencyeyecare/a/Sports_Injuries.htm, December 27, 2013.

Most Common Types of Eye Injuries

The most common types of eye injuries that can result from sports injuries are blunt injuries, corneal abrasions and penetrating injuries.

Blunt injuries: Blunt injuries occur when the eye is suddenly compressed by impact from an object. Blunt injuries, often caused by tennis balls, racquets, fists or elbows, sometimes cause a black eye or hyphema (bleeding in front of the eye). More serious blunt injuries often break bones near the eye, and may sometimes seriously damage important eye structures and/or lead to vision loss.

Corneal abrasions: Corneal abrasions are painful scrapes on the outside of the eye, or the cornea. Most corneal abrasions eventually heal on their

own, but a doctor can best assess the extent of the abrasion, and may prescribe medication to help control the pain. The most common cause of a sports-related corneal abrasion is being poked in the eye by a finger.

- + Penetrating injuries: Penetrating injuries are caused by a foreign object piercing the eye. Penetrating injuries are very serious, and often result in severe damage to the eye. These injuries often occur when eyeglasses break while they are being worn. Penetrating injuries must be treated quickly in order to preserve vision.⁴
- Pain when looking up and/or down, or difficulty seeing;
- Tenderness;
- Sunken eve;
- Double vision;
- Severe eyelid and facial swelling;
- Difficulty tracking;
- Symptoms of an Eye Injury
- The eye has an unusual pupil size or shape;
- Blood in the clear part of the eye;
- Numbness of the upper cheek and gum; and/or
- Severe redness around the white part of the eye.

What to do if a **Sports-Related Eye Injury** Occurs

If a child sustains an eye injury, it is recommended that he/she receive immediate treatment from a licensed HCP (e.g., eye doctor) to reduce the risk of serious damage, including blindness. It is also recommended that the child, along with his/her parent or guardian, seek guidance from the HCP regarding the appropriate amount of time to wait before returning to sports competition or practice after sustaining an eye injury. The school nurse and the child's teachers should also be notified when a child sustains an eye injury. A parent or guardian should also provide the school nurse with a physician's note detailing the nature of the eye injury, any diagnosis, medical orders for

the return to school, as well as any prescription(s) and/or treatment(s) necessary to promote healing, and the safe resumption of normal activities, including sports and recreational activities.

> According to the American Family Physician Journal, there are several guidelines that should be followed when students return to play after sustaining an eye injury. For

Return to Play and Sports

example, students who have sustained significant ocular injury should receive a full examination and clearance by an ophthalmologist or optometrist. In addition, students should not return to play until the period of time recommended by their HCP has elapsed. For more

minor eye injuries, the athletic trainer may determine that it is safe for a student to resume play based on the nature of the injury, and how the

student feels. No matter what degree of eye injury is sustained, it is recommended that students wear protective eyewear when returning to play and immediately report any concerns with their vision

to their coach and/or the athletic trainer.

Additional information on eye safety can be found at http://isee.nei.nih.gov and http://www.nei.nih.gov/sports.

⁴Bedinghaus, Troy, O.D., Sports Eye Injuries, http://vision.about.com/od/emergencyeyecare/a/Sports_Injuries.htm, December 27, 2013.

Signs or

Website Resources

- Sudden Death in Athletes http://tinyurl.com/m2gjmvq
- Hypertrophic Cardiomyopathy Association www.4hcm.org
- American Heart Association www.heart.org

Collaborating Agencies:

American Academy of Pediatrics New Jersey Chapter 3836 Quakerbridge Road, Suite 108 Hamilton, NJ 08619 Hamilton, VJ 08619 (p) 609-842-0014 (f) 609-842-0015



www.aapnj.org

American Heart Association 1 Union Street, Suite 301 Robbinsville, NJ, 08691 (p) 609-208-0020 www.heart.org

New Jersey Department of Education

PO Box 500 Trenton, NJ 08625-0500 (p) 609-292-5935 www.state.nj.us/education/

New Jersey Department of Health P. O. Box 360

r. 0. 80x 300 Trenton, NJ 08625-0360 (p) 609-292-7837 www.state.nj.us/health

NJ Health

Lead Author: American Academy of Pediatrics, New Jersey Chapter

Written by: Initial draft by Sushma Raman Hebbar, MD & Stephen G. Rice, MD PhD

Additional Reviewers: NJ Department of Education, NJ Department of Health and Senior Services, American Heart Association/New Jersey Chapter, NJ Academy of Family Practice, Pediatric Cardiologists, New Jersey State School Nurses

Revised 2014: Nancy Curry, EdM; Christene DeWitt-Parker, MSN, CSN, RN; Lakota Kruse, MD, MPH; Susan Martz, EdM; Stephen G. Rice, MD; Jeffrey Rosenberg, MD, Louis Teichholz, MD; Perry Weinstock, MD

SUDDEN CARDIAC DEATH

ATHLETES

The Basic Facts on Sudden Cardiac Death in Young Athletes



STATE OF NEW JERSEY DEPARTMENT OF EDUCATION

American Academy of Pediatrics





Substituting the set of the set o

What is sudden cardiac deatl in the young athlete?

Sudden cardiac death is the result of an unexpected failure of proper heart function, usually (about 60% of the time) during or immediately after exercise without trauma. Since the heart stops pumping adequately, the athlete quickly collapses, loses consciousness, and ultimately dies unless normal heart rhythm is restored using an automated external defibrillator (AED).

How common is sudden death in young athletes?

Sudden cardiac death in young athletes is very rare. About 100 such deaths are reported in the United States per year. The chance of sudden death occurring to any individual high school athlete is about one in 200,000 per year.

Sudden cardiac death is more common: in males than in females; in football and basketball than in other sports; and in African-Americans than in other races and ethnic groups.

What are the most common causes?

Research suggests that the main cause is a loss of proper heart rhythm, causing the heart to quiver instead of pumping blood to the brain and body. This is called ventricular fibrillation (ven-TRICK-you-lar fibroo-LAY-shun). The problem is usually caused by one of several cardiovascular abnormalities and electrical diseases of the heart that go unnoticed in healthy-appearing athletes.

The most common cause of sudden death in an athlete is hypertrophic cardiomyopathy (hi-per-TRO-fic CAR- dee-oh-my-OP-a-thee) also called HCM. HCM is a disease of the heart, with abnormal thickening of the heart muscle, which can cause serious heart rhythm problems and blockages to blood flow. This genetic disease runs in families and usually develops gradually over many years.

The second most likely cause is congenital (con-JEN-it-al) (i.e., present from birth)

abnormalities of the coronary arteries. This means that these blood vessels are connected to the main blood vessel of the heart in an abnormal way. This differs from blockages that may occur when people get older (commonly called "coronary artery disease," which may lead to a heart attack).

Other diseases of the heart that can lead to sudden death in young people include:

- Myocarditis (my-oh-car-DIE-tis), an acute inflammation of the heart muscle (usually due to a virus).
- Dilated cardiomyopathy, an enlargement of the heart for unknown reasons.
- Long QT syndrome and other electrical abnormalities of the heart which cause abnormal fast heart rhythms that can also run in families.
- Marfan syndrome, an inherited disorder that affects heart valves, walls of major arteries, eyes and the skeleton. It is generally seen in unusually tall athletes, especially if being tall is not common in other family members.

Are there warning signs to watch for?

In more than a third of these sudden cardiac deaths, there were warning signs that were not reported or taken seriously. Warning signs are:

- Fainting, a seizure or convulsions during physical activity;
- Fainting or a seizure from emotional excitement, emotional distress or being startled;
- Dizziness or lightheadedness, especially during exertion;
- Chest pains, at rest or during exertion;
- Palpitations awareness of the heart beating unusually (skipping, irregular or extra beats) during athletics or during cool down periods after athletic participation;
- Fatigue or tiring more quickly than peers; or
- Being unable to keep up with friends due to shortness of breath (labored breathing).

SUDDEN CARDIAC DEATH IN YOUNG ATHLETES

What are the current recommendations for screening young athletes?

- New Jersey requires all school athletes to be examined by their primary care physician ("medical home") or school physician at least once per year. The New Jersey Department of Education requires use of the specific Preparticipation Physical Examination Form (PPE).
- This process begins with the parents and student-athletes answering questions about
- symptoms during exercise (such as chest pain, dizziness, fainting, palpitations or shortness of breath); and questions about family health history.

The primary healthcare provider needs to know if any family member died suddenly during physical activity or during a seizure. They also need to know if anyone in the family under the age of 50 had an unexplained sudden death such as drowning or car accidents. This information must be provided annually for each exam because it is so essential to identify those at risk for sudden cardiac death.

The required physical exam includes measurement of blood pressure and a careful listening examination of the heart, especially for murmurs and rhythm abnormalities. If there are no warning signs reported on the health history and no abnormalities discovered on exam, no further evaluation or testing is recommended.

Are there options privately available to screen for cardiac conditions?

Technology-based screening programs including a 12-lead electrocardiogram (ECG) and echocardiogram (ECHO) are noninvasive and painless options parents may consider in addition to the required

PPE. However, these procedures may be expensive and are not currently advised by the American Academy of Pediatrics and the American College of Cardiology unless the PPE reveals an indication for these tests. In addition to the expense, other limitations of technology-based tests include the possibility of "false positives" which leads to unnecessary stress for the student and parent or guardian as well as unnecessary restriction from athletic participation.

The United States Department of Health and Human Services offers risk assessment options under the Surgeon General's Family History Initiative available at http://www.hhs.gov/familyhistory/Index.html.

When should a student athlete see a heart specialist?

If the primary healthcare provider or school physician has concerns, a referral to a child heart specialist, a pediatric cardiologist, is recommended. This specialist will perform a more thorough evaluation, including an electrocardiogram (ECG), which is a graph of the electrical activity of the heart. An echocardiogram, which is an ultrasound test to allow for direct visualization of the heart structure, will likely also be done. The specialist may also order a treadmill exercise test and a monitor to enable a longer recording of the heart rhythm. None of the testing is invasive or uncomfortable.

Can sudden cardiac death be prevented just through proper screening?

A proper evaluation should find most, but not all, conditions that would cause sudden death in the athlete. This is because some diseases are difficult to uncover and may only develop later in life. Others can develop following a

normal screening evaluation, such as an infection of the heart muscle from a virus.

This is why screening evaluations and a review of the family health history need to be performed on a yearly basis by the athlete's primary healthcare provider. With proper screening and evaluation, most cases can be identified and prevented.

Why have an AED on site during sporting events?

The only effective treatment for ventricular fibrillation is immediate use of an automated external defibrillator (AED). An AED can restore the heart back into a normal rhythm. An AED is also life-saving for ventricular fibrillation caused by a blow to the chest over the heart (commotio cordis).

N.J.S.A. 18A:40-41a through c, known as "Janet's Law," requires that at any schoolsponsored athletic event or team practice in New Jersey public and nonpublic schools including any of grades K through 12, the following must be available:

- An AED in an unlocked location on school property within a reasonable proximity to the other of the second s
- A team coach, licensed athletic trainer, or other designated staff member if there is no coach or licensed athletic trainer present
- coach or licensed athletic trainer present, certified in cardiopulmonary resuscitation (CPR) and the use of the AED; or
- A State-certified emergency services provider or other certified first responder.

The American Academy of Pediatrics recommends the AED should be placed in central location that is accessible and ideally no more than a 1 to $1/_2$ minute walk from any location and that a call is made to activate 911 emergency system while the AED is being retrieved.

State of New Jersey DEPARTMENT OF EDUCATION

Sudden Cardiac Death Pamphlet

Sign-Off Sheet

Name of School District:_____

Name of Local School: ____

I/We acknowledge that we received and reviewed the Sudden Cardiac Death in Young Athletes pamphlet.

.

Student Signature:

Parent or Guardian Signature:

Date:____

New Jersey Department of Education 2014: pursuant to the Scholastic Student-Athlet Safety Act, P.L. 2013, c71